Page 1 of 2 School Asthma Action Plan

				Dist. D. 4				····	
Name				Birth Date Address		Address			
Emergency Contact				Phone			Cell		
Triggers	⊒ Mold/Pollens	☐ Animals ☐ Co	lds 🔱 Du	st 🗀 Exe	ercise	⊒ Smoke	☐ Weather	☐ Fragrance	
Green Zone: Doing Well *Breathing is good •No cough or wheeze •Can work and play •Sleeps all night *No early warning signs •Peak Flow Meter if used: 80-100% of personal best									
School Action: Follow actions in marked boxes below for exercise induced asthma									
□ Medication Before Exercise □ Medication Before Recess □ Use routinely every hours									
Medication with spacer: □ Albuterol □ Ventolin □ Proventil □ Xopenex									
Medication without spacer: □ Maxair Autohaler Dose: puffs When: 10-15 minutes before listed activity Start Date: School Year Stop Date: School Year									
Source 14 treath 10 10 minutes octobe usten animals State Nater School 1681. State School 1681.									
	one: Getting W	 Cough, wheere, chest tight - Problems working/ playing Early warning signs - Shortness of breath Peak Flow Meter if used: 50 to 60% of personal best 							
School Actions: Follow actions in marked boxes below									
Take Quick-Relief Medication			How Much (Dose)		When		Start Date	Stop Date	
MDI with Spacer: □ Albutero: □ Ventolin □ Proventi: □ Xopenex			puffs		Student symptoi	report of ns	School Year	School Year	
Without spacer: Maxair Autohaler									
Nebutizer: □ Albutero: □ Ventolin □ Proventil □Xopenex			u			report of ns	School Year	School Year	
□ If symptoms improve after 10-15 minutes: Return to normal activity									
☐ If symptoms do not improve after 10-15 minutes: Give quick relief medication again and call parents									
		r the second 10-15					-		
□ If symptoms do not improve after the medication is repeated: Call EMS (911), School RN and parents □ If symptoms get worse at anytime: Call EMS (911), School RN and Parents									
		anytime: Call EMS k relief medications					e School RN ar	nd Parents	
Red Zone: Medical Alert (severe trouble breathing) Cannot stop coug Getting worse, in The skin between Lips or fingernals			stead of better • Trouble walking or talking from shortness of breath the ribs and above the collarbone pulls in or retracts						
School Actions: 1. Call EMS (911) IMMEDIATELY 2. GIVE QUICK-RELIEVER MEDICATION AND CONTINUE EVERY 15 MINUTES UNTIL EMS (911) ARRIVES 3. Call School RN and Parents									
Take Quick-Relief Medications			How Mu	ch (Dose)	When		Start Date	Stop Date	
MDI with Spacer: □ Albuterol □ Ventolin □ Proventil □ Xopenex Without spacer: □ Maxair Autohaler				puffs		t report of ervation of oms.	School Year	School Year	
Nebulizer: □ Albuterol □ Ventolin □ Proventil □Xopenex				llait ta		t report of ervation of oms	School Year	School Year	
Heath Co.	Daniella a N	_		Unit Dose					
Heath Care Provider Name: Phone: FAX#:									
Health Care Provider Sign: Date:									

Page 2 of 2

Student Name:	Birth Date:					
Address: School Asthma Action Plan (page two)						
School Asthma Action Plan (page two)						
Metered Dose Inhaler (MDI) Instructions						
1. Store at room temperature.						
2. Shake the MDI for 5 seconds before each						
 Prime the MDI before the first use or when product's patient information sheet for MD usually involves pressing down on the medication one or more puff of medication. Discarding contains the labeled amount of medication Keep track of metered inhalation puffs use number of metered inhalation puffs available metered inhaled puffs available is listed on 	OI specific priming instructions. Priming dication canister to discard into the air g puffs makes sure the next puff inhaled. d. Subtract the number used from the pole listed on the label. The number of					
There are usually 200 puffs in an MDI. 4. Ask family for a new MDI when all labeled	d metered inhalation puffs are used.					
MDI and Aerosol Solution Potential Adverse Reaction: Headache, shakiness, fast heart rate, nausea. Call parent with 1) student report of symptoms that interfere with schoolwork or activity 2) increase in side effects 3) frequent usage (2 times a day for 3 days). We have instructed the patient and family in the proper use of the quick-relief medications. It is my professional opinion that the student: should be allowed to carry and self administer the inhaled medication. The medication should be stored and administered by designated school personnel.						
Provider Signature	Date					
**************************************	************					
I give permission for my child to receive medication at school in keeping with Section I above according to the school district policy and as instructed by the physician and agree to 1) Assume responsibility for safe delivery of the medication in its original container to the school, 2) Have a new form completed by the doctor if medication or dosage is changed, 3) Notify the school of changes in health care provider. I release from liability, and in addition agree to indemnify, all school employees, the Board of Education and Akron Children's Hospital School Health Services for damages or injury resulting from the use, misuse or nonuse of such medication except as such Board, School Health Services or its employees are grossly negligent or engage in wanton or reckless misconduct. I further agree to submit a revised statement signed by the physician who has prescribed the medication described in Section I in the event that I become aware that any of the above information has changed. I have read and understand the policy of the School Board for administration of medication and affirm that this request entails special circumstances justifying an exception from the usual administration of medication at school-by-school personnel.						
Parent/Guardian Signature:	Date:					
Daytime Phone: THIS FORM EXPIRES AT THE END OF THE SCHOOL YEAR						